

On the Consistency of Twisted Gauge Theory ¹

Stefan Giller[†]
Cezary Gonera[‡]
Piotr Kosiński[‡]
Paweł Maślanka[‡]

[†]Jan Dlugosz Academy in Czestochowa
Institute of Physics
ul. Armii Krajowej 13/15, 42-200 Czestochowa, Poland
e-mail: sgiller@uni.lodz.pl

[‡]Theoretical Physics Department II, University of Łódź,
Pomorska 149/153, 90-236 Łódź, Poland
e-mail: cgonera@uni.lodz.pl
pkosinsk@uni.lodz.pl
pmaslan@uni.lodz.pl

Abstract

It is argued that the twisted gauge theory is consistent provided it exhibits also the standard noncommutative gauge symmetry.

¹Supported by the grant no 1P03B02128 of the Polish Ministry of Science

In the recent years the noncommutative field theories [1, 2, 3] have gained much attention. This is in part due to the fact that they emerge as a specific limit of string theory and, being much simpler than the original string theory, yet keep some of its properties [4, 5]. Moreover, they can be viewed as an attempt to describe the space-time at very short distances where the very idea of space-time as a smooth manifold may lack any meaning.

Due to the crucial role played by gauge theories it is natural to ask whether the idea of gauge symmetry can be generalized to the case of the field theory on noncommutative space-time. When trying to extend in this direction the notion of gauge transformations one is faced with some serious problems because for many classical groups the modification consisting in replacing the product with star product leads to the transformations which do not close to form a group. These problems lead to some no-go theorems restricting the group and representing content of noncommutative gauge theories [6]. A number of more or less sophisticated proposals has been made to avoid the troubles with the idea of the localization of symmetry on noncommutative space-time [7, 8, 9].

The most flexible one seems to be the approach based on the idea of twisted gauge symmetry [10, 11, 12, 13]. Putting aside all formal details one can summarize the notion of twisted gauge transformations as follows: in order to compute the gauge variation of star-product monomial in fields and their derivatives one applies the Leibnitz rule but always keeping the gauge parameter function leftmost. It is easily seen that in this case any gauge group can be closed and the twisted gauge invariance imposes rather mild restrictions on the form of the action. The idea of twisted gauge transformations seems to provide a general method of introducing the gauge symmetry for field theories on noncommutative space-time. However, according to Milton Friedman, there are no lunches for free. This observation applies here as well. First it has been argued the the approach based on twisted gauge transformations is in contradiction with the very concept of gauge fields [14].

Another serious arguments are given in the paper of Alvarez-Gaume *et al.* [15]. They noticed that the modified Leibnitz rule (due to the fact that the gauge parameter function is kept leftmost) means that the twisted symmetry is not a physical symmetry in the usual sense and it is not clear whether the Noether theorem can be derived. For the case of twisted $U(N)$ gauge theory the action is also invariant under the standard noncommutative $U(N)$ symmetry and this could be the reason for the existence of conserved current. This observation may be linked with the one made in Refs. [11, 12]. It has been observed [11, 12] that, in spite of the fact that one can gauge any group, when writting the invariant action one must admit that the gauge potential takes values in larger algebra than the Lie algebra of the gauge group. This is not only because the field strength takes necessarily values in enveloping algebra but is also somehow necessary for the consistency of field equations [12]. In fact, it appears that in all examples considered so far the range of values of the gauge potential is such that the action invariant under twisted gauge transformations is also invariant under standard noncommutative gauge symmetry; the latter enforces the consistency of current conservation and field equations.

In the present note we argue that this is generally the case. Namely, if one considers the theory invariant under twisted gauge symmetry then, in general, the current conservation implies some constraints which are not automatically fulfilled by virtue of field equations; more specifically, these constraints follow by taking the divergence of field equtaions. If one demands no such constraints exist, the range of gauge potentials must be enlarged so that the action exhibits also standard noncommutative gauge symmetry. Therefore, the twisted gauge theory, if consistent, posses a custodial standard symmetry enforcing current conservation.

For gauge theories on commutative space-time the local gauge invariance allows to derive the set of identities [16] implying that the divergencies of Euler-Lagrange expressions vanish by virtue of field equations. We explain why this is also the case for noncommutative gauge symmetry but not for the twisted one. Roughly speaking, this is because the variation of action under standard noncommutative gauge symmetry is computed in the same way as the variation leading to the field equations. More detailed discussion is given in the final part of the paper.

We start with the following general example.

Let G be some compact Lie group with the Lie algebra g . We want to construct the gauge theory on noncommutative space-time based on twisted gauge symmetry with G playing the role of gauge group. To this end we select some N -dimensional unitary irreducible representation of G (and the corresponding irreducible representation of g , denoted also by g) and assume that the gauge potential takes its values in g . The Lie algebra g is a subalgebra of $u(N)$. The latter viewed as linear space is equipped with the scalar product $(A, B) \equiv \text{Tr}(AB)$ invariant under the adjoint action of $U(N)$. Let $g_\perp$ be the orthogonal complement of g in $u(N)$ and let B_i , $i = 1, \dots, a$, be an orthonormal bases in $g_\perp$. The subspace g can be now characterized by the condition: $A \in g \iff \text{Tr}(AB_i) = 0$, $i = 1, \dots, a$.

The gauge theory with twisted G -symmetry is now formulated with help of Lagrange multiplier method. To this end we assume that the gauge potential A_μ *a priori* takes its values in $U(N)$ and consider the action:

$$S = -\frac{1}{4} \int d^4x \text{Tr}(F_{\mu\nu} * F_{\mu\nu}) + \int d^4x \rho_i^\mu \text{Tr}(A_\mu B_i) \quad (1)$$

where

$$F_{\mu\nu} \equiv \partial_\mu A_\nu - \partial_\nu A_\mu - i[A_\mu, A_\nu]_* \quad (2)$$

while $\rho_i^\mu(x)$ are the Lagrange multipliers.

Eq. (1) implies the following equations of motion

$$\begin{aligned} \text{Tr}(A_\mu B_i) &= 0 \\ \partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_* + \rho_i^\nu B_i &= 0 \end{aligned} \quad (3)$$

Eliminating the Lagrange multipliers one obtains

$$\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_* - \text{Tr}(B_i(\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_*))B_i = 0 \quad (4)$$

When projected on g the above equation yields standard Yang-Mills equation

$$\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_*|_g = 0 \quad (5)$$

However, one must keep in mind that even if the gauge potential takes its values in g this is no longer the case for the field strength tensor $F_{\mu\nu}$. Therefore, $\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_*$ can have a nonvanishing component in $g_\perp$. In fact, eq.(4) is just the statement that $\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_* \in g_\perp$.

In the case of commutative space-time $\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_* \in g$ by construction so the field equations imply $\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_* = 0$. In the standard (i.e. not twisted) approach to gauge theory on noncommutative space-time g must basically be $u(N)$ and $g_\perp = 0$; again one arrives at standard form of field equations.

Having obtained the field equations we can ask whether they are consistent with the requirement of current conservation. This seems to be by far not obvious. In fact, taking the divergence of eq.(4) results in the following additional condition:

$$Tr(\Gamma[A_\nu, Tr(B_i(\partial_\mu F^{\mu\nu} - i[A_\mu, F^{\mu\nu}]_*)B_i)_*]) = 0 \quad (6)$$

for any $\Gamma \in g$. We would like to know under which circumstances eq.(6) does not provide further constraints on gauge fields. Let us note that the star commutator under the outer trace consists of two pieces, one proportional to the matrix commutator and the second one involving matrix anticommutator. Now by virtue of field equations (2) A_μ belongs to g . It is straightforward to check that $[\Gamma', B_i] \in g$ for any $\Gamma' \in g$. Indeed, if $U \in G$ then $U^+ \Gamma' U \in g$ and $Tr(\Gamma' U B_i U^+) = Tr(U^+ \Gamma' U B_i) = 0$ so that $U B_i U^+ \in g_\perp$. Therefore, the part involving commutator gives no constraint. The piece containing the anticommutator does not product any constraint only provided $\{\Gamma', B_i\} \in g_\perp$ for $\Gamma' \in g$. This together with $[\Gamma', B_i] \in g_\perp$ implies that both $\Gamma' B_i \in g_\perp$ and $B_i \Gamma' \in g_\perp$ so that $\Sigma B_i \in g_\perp$ and $B_i \Sigma \in g_\perp$ for each Σ belonging to the enveloping algebra of g . By assumption g is irreducible; the Burnside theorem implies then that B_i span two-sided ideal in the algebra of all $N \times N$ matrices. However, by virtue of Wedderburn theorem such an ideal, if proper (only then there is some gauge symmetry at all) must be zero. Then $g = u(N)$ and we are dealing with the theory exhibiting standard noncommutative gauge symmetry.

Consider some more specific examples of the situation described above. In Ref[11] the $SU(2)$ twisted gauge theory is considered. However, it is admitted from the very beginning, that the gauge field takes its value in $U(2)$ algebra and that the resulting theory is invariant under the standard $U(2)$ noncommutative gauge symmetry which guarantees the existence of the conserved current. On the other hand, if we assume that A_μ belongs to the $SU(2)$ algebra only the second equation (4.29) of Ref.[11] survives and, additionally, $B_\mu = 0$. One easily checks that by taking the divergence of this equation one obtains a new constraint of the form (6).

As a second example take $G = SO(N)$ in the defining representation. Then g is the set of imaginary antisymmetric $N \times N$ matrices while $g_\perp$ - the set of the real symmetric ones. Again one can write the Lagrangian containing gauge fields taking their values in $SO(N)$. Taking the divergence of the resulting field equations and separating symmetric and antisymmetric parts one arrives at the constraint of the general form (6).

Finally, let us consider the simple example of gauge theory with matter fields [12]. The gauge group is $U(1)$ and the matter fields form the multiplet transforming as follows

$$\delta_\alpha \Psi = i\alpha \mathbf{Q} \Psi, \quad \mathbf{Q} = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} \quad (7)$$

The gauge field takes its values in the algebra spanned by $\mathbf{I}$ and $\mathbf{Q}$

$$\mathbf{A}_\mu = A_\mu \mathbf{I} + B_\mu \mathbf{Q} \quad (8)$$

and the Lagrangian reads

$$L = -\frac{1}{4} Tr(\mathbf{F}^{\mu\nu} * \mathbf{F}_{\mu\nu}) + \bar{\Psi} * \gamma^\mu (i\partial_\mu + A_\mu) \Psi - m \bar{\Psi} * \Psi \quad (9)$$

where

$$\mathbf{F}_{\mu\nu} \equiv \partial_\mu \mathbf{A}_\nu - \partial_\nu \mathbf{A}_\mu - i[\mathbf{A}_\mu, \mathbf{A}_\nu]_* \quad (10)$$

In order to reveal the content of the theory we define two projectors

$$\mathbf{P}_\pm \equiv \frac{1}{2}(\mathbf{I} \pm \mathbf{Q}) \quad (11)$$

and the corresponding decomposition of the gauge field

$$\mathbf{A}_\mu = A_\mu^+ \mathbf{P}_+ + A_\mu^- \mathbf{P}_-, \quad A_\mu^\pm \equiv A_\mu \pm B_\mu \quad (12)$$

and the matter field

$$\Psi_\pm \equiv \mathbf{P}_\pm \Psi, \quad \Psi = \Psi_+ + \Psi_- \quad (13)$$

Then

$$\mathbf{F}_{\mu\nu} = F_{\mu\nu}^+ \mathbf{P}_+ + F_{\mu\nu}^- \mathbf{P}_- \quad (14)$$

with

$$F_{\mu\nu}^\pm \equiv \partial_\mu A_\nu^\pm - \partial_\nu A_\mu^\pm - i[A_\mu^\pm, A_\nu^\pm]_*$$

$$(\partial_\mu - iA_\mu^*)\Psi = (\partial_\mu - iA_\mu^+*)\Psi_+ + (\partial_\mu - iA_\mu^-*)\Psi_- \quad (15)$$

Therefore, the Lagrangian splits into two independent parts

$$L = -\frac{1}{4}Tr(F^{+\mu\nu} * F_{\mu\nu}^+) + \bar{\Psi}_+ * \gamma^\mu (i\partial_\mu + A_\mu^+*)\Psi_+ - m\bar{\Psi}_+ * \Psi_+ -$$

$$\frac{1}{4}Tr(F^{-\mu\nu} * F_{\mu\nu}^-) + \bar{\Psi}_- * \gamma^\mu (i\partial_\mu + A_\mu^-*)\Psi_- - m\bar{\Psi}_- * \Psi_- \quad (16)$$

The "±" degrees of freedom decouple and both pieces exhibit standard (noncommutative) $U(1)$ symmetry.

The important point here is that the gauge potential is a matrix acting in representation space of matter fields. However, consistency (i.e. the absence of further constraints) demands that A_μ takes its values in the enveloping algebra of the Lie algebra under consideration. Therefore, the gauge potential is the set of two independent fields and the theory describes two independent standard noncommutative $U(1)$ -symmetric pieces.

On the other hand, if we insist on using the gauge potential belonging to the Lie algebra $U(1)$ (i.e. consisting of one field) then either all fields have the same charge (i.e. the enveloping algebra coincides with the Lie algebra of $U(1)$) or there is additional constraint[12].

In all the above examples in order to avoid additional constraints following from the field equations one has to extend the definition of gauge potential in such a way that the resulting theory exhibits the standard noncommutative gauge symmetry. It seems that just the very existence of the latter enforces consistency.

In order to put more light on the problem let us remind the second Noether theorem in the commutative case[16]. Consider the field theory described by the Lagrangian $L(\Phi, \partial_m u\Phi)$ carrying the following local symmetry

$$\delta\Phi^i(x) = a_k^i(\Phi; x)\epsilon^k(x) + b_k^{i\mu}(\phi; x)\partial_\mu\epsilon^k(x) \quad (17)$$

Then the invariance of L implies the following identities

$$\begin{aligned}
a_k^i \frac{\delta L}{\delta \Phi^i} + \partial_\mu \left(a_k^i \frac{\partial L}{\partial (\partial_\mu \Phi^i)} \right) &\equiv 0 \\
b_k^{i\mu} \frac{\delta L}{\delta \Phi^i} + a_k^i \frac{\partial L}{\partial (\partial_\mu \Phi^i)} + \partial_\nu \left(b_k^{i\mu} \frac{\partial L}{\partial (\partial_\mu \Phi^i)} \right) &\equiv 0 \\
b_k^{i\nu} \frac{\partial L}{\partial (\partial_\mu \Phi^i)} + b_k^{i\mu} \frac{\partial L}{\partial (\partial_\nu \Phi^i)} &\equiv 0
\end{aligned} \tag{18}$$

where

$$\frac{\delta L}{\delta \Phi^i} \equiv \frac{\partial L}{\partial \Phi^i} - \partial_\mu \left(\frac{\partial L}{\partial (\partial_\mu \Phi^i)} \right)$$

is the Euler – Lagrange expression.

The first identity expresses the conservation of the current

$$j_k^\mu \equiv a_k^i \frac{\partial L}{\partial (\partial_\mu \Phi^i)} \tag{19}$$

and is valid even if L exhibits merely global symmetry. The remaining identities hold only for local symmetries. They imply that the field equations are in fact dependent. Indeed, one easily derives the following Bianchi identities

$$a_k^i \frac{\delta L}{\delta \Phi^i} - \partial_\mu \left(b_k^{i\mu} \frac{\delta L}{\delta \Phi^i} \right) = 0 \tag{20}$$

In the particular case of pure gauge theory eqs.(14),(18) take the following form (with c_{bc}^a being the structure constants)

$$\begin{aligned}
c_{bc}^a \left(A_\mu^c \left(\frac{\partial L}{\partial A_\mu^a} - \partial_\nu \left(\frac{\partial L}{\partial (\partial_\nu A_\mu^a)} \right) \right) + \partial_\nu \left(A_\mu^c \frac{\partial L}{\partial (\partial_\nu A_\mu^a)} \right) \right) &= 0 \\
\frac{\partial L}{\partial A_\mu^a} - \partial_\nu \left(\frac{\partial L}{\partial (\partial_\nu A_\mu^a)} \right) - c_{ac}^b A_\nu^c \frac{\partial L}{\partial (\partial_\mu A_\nu^b)} + \partial_\nu \left(\frac{\partial L}{\partial (\partial_\nu A_\mu^a)} \right) &= 0 \\
\frac{\partial L}{\partial (\partial_\nu A_\mu^a)} + \frac{\partial L}{\partial (\partial_\mu A_\nu^a)} &= 0
\end{aligned} \tag{21}$$

while (20) corresponds to

$$c_{bc}^a A_\mu^c \frac{\delta L}{\delta A_\mu^a} - \partial_\mu \left(\frac{\delta L}{\delta A_\mu^b} \right) = 0 \tag{22}$$

We see from eq.(22) that no new relation (constraint) follows by taking the divergence of field equations.

In the case noncommutative field theories gauge symmetry also implies the set of identities. Let us consider first the standard noncommutative gauge theory. In order to answer the question whether there exists some counterpart of Noether identities (21) let us recall the way the field equations are obtained from variational principle. One computes the variation of the

Langrangian by applying Leibnitz rule to each monomial entering the latter and making cyclic permutations to bring the field variation say leftmost. Consequently, the field equations can be written in standard form provided the derivatives $\frac{\partial}{\partial A_\mu^a}, \frac{\partial}{\partial(\partial_\nu A_\mu^a)}$ are defined as follows: in order to apply the relevant derivatives to some monomial in gauge fields and their derivatives one uses the Leibnitz rule and then makes in each term the cyclic permutation to bring the derivative leftmost. On the other hand, similar situation is encountered when considering the standard noncommutative gauge symmetry. Again, one considers the variation of the Lagrangian implied, via the Leibnitz rule, by the variation of gauge fields under the gauge transformation. The variation of action is obtained by varying the Lagrangian under the integral so one can make cyclic transposition to bring the gauge parameter leftmost. The result is that the coefficients behind gauge parameter and its first and second derivatives vanish. In this way we obtain a set of identities in which derivatives $\frac{\partial}{\partial A_\mu^a}, \frac{\partial}{\partial(\partial_\nu A_\mu^a)}$ have the same meaning as in the case of field equations. These identities have therefore very similar form to their commutative counterparts. In particular, they also imply that no further constraint follows by taking the divergence of field equations.

Let us now turn to twisted gauge symmetry. The formal definition of twisted gauge transformation can be summarized as follows: the variation of arbitrary monomial in gauge fields and their derivatives is calculating according to the Leibnitz rule but with the gauge parameter put leftmost (no cyclic permutation). The derivatives $\frac{\partial}{\partial A_\mu^a}, \frac{\partial}{\partial(\partial_\nu A_\mu^a)}$ entering the identities expressing the invariance under the twisted gauge symmetry are therefore defined by Leibnitz rule (without cyclic permutation). In spite of the fact that they appear in the combination resembling Euler – Lagrange expressions such combinations do not vanish by virtue of the field equations due to the difference in the definition of derivatives. As a result, the modified Noether identities do not guarantee the consistency of field equations. This seems to make the notion of twisted gauge symmetry less interesting.

The structure of twisted gauge theories has been also analysed in two recent papers [17], [18]

References

- [1] Douglas M.R., Nekrasov N.A., *Rev. Mod. Phys.* **73** (2001) 977
- [2] Szabo R.J., *Phys. Rep.* **378** (2003) 207
- [3] Schaposhnik F.A., *Three lectures on noncommutative field theories*, hep-th/0408132
- [4] Connes A., Douglas M.R., Schwarz A., *JHEP* **02** (1998) 003
- [5] Douglas M.R., Hull C., *JHEP* **02** (1998) 008
- [6] Chaichian M., Presnajder P., Sheikh-Jabbari M.M., Tureanu A., *Phys. Lett.* **B526** (2002) 132
- [7] Bonora L., Schnabl M., Sheikh-Jabbari M.M., Tomasiello A., *Nucl. Phys.* **B589** (2000) 461
- [8] Bars J., Sheikh-Jabbari M.M., Vasiliev M.A., *Phys. Rev.* **D64** (2001) 086004
- [9] Jurco J., Schraml S., Schupp P., Wess J., *Eur. Phys. J.* **C17** (2000) 521

- [10] Vassilevich D.V., *Mod. Phys. Lett.* **A21** (2006) 1279
- [11] Aschieri P., Dmitrijevich M., Meyer F., Schraml S., Wess J., *Lett. Math. Phys.* **78** (2006) 61
- [12] Wess J., *Deformed Gauge Theories*, (hep-th/0608135)
- [13] Garcia-Bondia J.M., Lizzi F., Ruiz Ruiz F., Vitale P., *Phys. Rev.* **D74** (2006) 025015
- [14] Chaichian M., Tureanu A., *Phys. Lett.* **B637** (2006) 199
- [15] Alvarez-Gaume L., Meyer F., Vazquez-Moro M.A., *Nucl. Phys.* **B753** (2006) 92
- [16] Konopleva N., Popov V., *Kalibrovochnye polja* Atomizdat, Moscow 1980 (in Russian)
- [17] Banerjee R., Samanta S., *Gauge Generators, Transformations and Identities on a Non-commutative Space*, hep-th/0608214
- [18] Banerjee R., Samanta S., *Gauge Symmetries on Θ -Deformed Spaces*, hep-th/0611249